

FOR USE IN STORAGE SITES:

This concentrate can be used to treat grain and seed in warehouse bins and trucks, cargo ships, mills, bin hoppers, elevators and conveying equipment as a clean-up prior to using them for storage. In mills and elevators, all infested accumulations of grain should be removed from the bin hoppers. All storage areas and conveying equipment should be thoroughly cleaned by sweeping out the waste grain, cobwebs and other debris from the walls and rafters as well as on the floor and door frames with special attention to material lodged in the cracks and crevices. All of the debris should be removed and burned to kill eggs and insects that might be present.

For farms, particular attention should be given to cleaning up around the used feed and grain bags, grain residues from wagons, harvesting equipment and feed troughs. Newly harvested grain should not be placed in the same bin with carry-over grain and all carry-over grain stocks that are not treated with grain protectant should be fumigated. These cleaning operations should be done within two or three weeks before harvest.

After above sanitation measures have been employed, spray all areas prior to use for storage with 1 part to 29 parts water (1 quart with 7.5 gallons water) up to 1 part to 11 parts (1 quart with 3 gallons water). Apply at the rate of one gallon per 750 square feet on walls, floors, ceilings and partition boards of bins, paying particular attention to forcing the spray into all cracks and crevices.

Monthly inspections should be made. If the top 2 or 3 inches are found to be infested, re-treat applying at the rate of 1 to 2 gallons of diluted material per 1,000 bushels of stored product.

FOR USE AS A LIVESTOCK AND POULTRY SPRAY:

1) To kill and repel horn flies, houseflies, mosquitoes and gnats, dilute at the rate of 1 to 2 fluid ounces per gallon of water and apply to wet the hair thoroughly with particular attention to topline, underline, flanks, withers and other infested areas. Repeat treatment at intervals of 5 to 12 days for small insect populations or as needed when flies are emerging in large numbers.

2) To kill and repel stable flies, horse flies, and deer flies, dilute at the rate of 2 to 3 fluid ounces per gallon of water and apply at a quart per adult animal to wet the hair thoroughly with particular attention to the legs, flanks, barrel, topline and other body areas commonly attacked by these flies. Repeat treatment each week as needed.

3) To kill face flies dilute at the rate of 2 fluid ounces per gallon of water and apply using spray which produces large wetting droplets. Apply to the face of the animal in the morning before releasing to pasture. Apply sufficiently to wet the face but not more than 1.5 ounces per animal. Repeat daily as needed.

4) For effective kill of biting and sucking lice on cattle, horses, sheep, goats and hogs, dilute at the rate of 1 quart with 75 gallons of water (1 fluid ounce with 2 gallons) and spray to thoroughly wet the hair of the animal including the head and brush of the tail. Repeat treatment in 10 days to kill newly hatched lice.

5) To kill poultry lice, using a dilution of 2 to 3 ounces of concentrate per gallon of water spray roosts, walls and nests or cages thoroughly. It is not necessary to remove poultry from the housing unit during treatment. This should be followed by spraying over the birds with a fine mist.

6) To kill bed bugs and mites on poultry and in poultry houses, dilute at the rate of 2 to 3 fluid ounces per gallon of water and spray crevices of roost poles, cracks in walls and cracks in nests where the bed bugs and mites hide, followed by spraying over the birds with a fine mist.

7) To kill sheep "tick" or ked, dilute at the rate of 1 to 2 fluid ounces per 4 gallons of water and thoroughly wet all portions of the body by dipping or by spraying with sufficient pressure and with a nozzle adjustment to give penetration of the wool. Treat at a rate sufficient to wet the animal.

8) To kill fleas and ticks on livestock and pets, and to obtain protection against reinfestation, dilute at the rate of 2 fluid ounces per gallon of water and wet the animal by dipping or spraying. For best results against fleas and ticks on dogs and cats the kennels and/or animal quarters and bedding should be treated.

FOR USE IN BARN, DAIRIES, MILKING PARLORS, MILKING ROOMS AND POULTRY HOUSES:

To kill flying insects including, but not limited to, Flies, Fruit Flies, Mosquitoes, Gnats, Wasps, Hornets and Small Flying Moths, dilute at the rate of 2 fluid ounces per gallon of water. Apply as a fog or fine mist (at approximately 2 ounces per 1,000 cubic feet of space), directing the nozzle for maximum coverage and above livestock and poultry toward the ceiling and upper corners of the area being treated. For best results, close doors and windows before spraying, and keep them closed for 10 to 15 minutes. Applicator should vacate the treated area and ventilate it prior to returning. Repeat application as necessary.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry area. Always store pesticides in the original container. Store away from food and pet food.

PESTICIDE DISPOSAL: To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill. **Triple rinse as follows (for containers of 5 gallons or less):** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Avoid contact with skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wear long-sleeved shirt and long pants, socks, shoes and chemical-resistant gloves (such as Barrier Laminate, Nitrile Rubber, Neoprene Rubber or Viton, Selection Category E). Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are made of barrier laminate, butyl rubber, nitrile rubber or viton. If you want more options, follow the instructions for category F on an EPA Chemical-resistance chart.

Applicators and other handlers who may be exposed to the dilute and/or concentrate through application or other tasks must wear:

- Long-sleeved shirt,
- Long pants, Chemical-resistant gloves, and
- Shoes plus socks; and
- Protective eyewear.

In addition to the above PPE, applicators using hand-held foggers in an enclosed area must wear a half-face, full-face or hood-style NIOSH-approved respirator with:

- A dust/mist filtering cartridge (MSHA/NIOSH approval number prefix TC-21C), or
- A canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G), or
- A cartridge or canister with any R, P, or HE filter.

See engineering controls for additional requirements.

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.



EPA Reg. No. 1021-1770
EPA Est. No. 1021-MN-2
0712-0613

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

IF SWALLOWED:

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

HOTLINE NUMBER: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For additional information on this pesticide product (including health concerns, medical emergencies or pesticide incidents), you may call 1-888-740-8712.

ENGINEERING CONTROLS

Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(6)].

Human flagging is prohibited. Flagging to support aerial application is limited to use of the Global Positioning System (GPS) or mechanical flaggers.

User Safety Recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic organisms, including fish and invertebrates. Drift and run-off may be hazardous to aquatic organisms in water adjacent to treated areas. This product may contaminate water through run-off. This product has a potential for run-off for several weeks after application. Poorly draining soils and soils with shallow water tables are more prone to produce run-off that contains this product.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

Except as specified in the directions for use, do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash-waters or rinsate.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.



Specimen Label



- Multi-purpose insecticide
- For use on growing crops, ornamentals, stored products, livestock and to kill fruit flies on harvested fruits and vegetables
- Used alone as a clean-up spray or a pre-harvest spray

ACTIVE INGREDIENTS:

Pyrethrins.....	6.00%
*Piperonyl butoxide, Technical.....	60.00%
OTHER INGREDIENTS.....	34.00%
	100.00%

*Equivalent to (butylcarbityl) (6-propylpiperonyl) ether and related compounds.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See pages 4 and 5 for first aid and precautionary statements.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS & PRECAUTIONS

APPLICATION RESTRICTIONS:

- Apply this product only as specified on this label.
- See separate directions and precautions for mosquito control applications.
- Do not contaminate food or feedstuffs.
- Except when applying this product as a wide-area mosquito adulticide, do not enter or allow others to enter until vapors, mists and aerosols have dispersed, and the area has been thoroughly ventilated.
- Except when applying to livestock and pets and when used as a wide area mosquito adulticide, do not apply this product in a way that will contact workers or other persons, either directly or through drift.
- Except when applying to livestock and pets and when used as a wide area mosquito adulticide, only protected handlers may be in the area during application.
- Do not remain in treated area. Exit area immediately and remain outside the treated area until aerosols, vapors and/or mists have dispersed.
- Remove or cover exposed food and drinking water before application.
- Remove or cover dishes, utensils, food processing equipment and food preparation surfaces, or wash them before use.
- For direct application to non-domestic animals/livestock: Do not apply more than 1 time per day.
- Do not make applications during rain.
- Do not water treated area to point of runoff.
- When used in dairy barns or facilities: Close milk bulk tank lids to prevent contamination from spray and from dead or falling insects. Remove or cover milking utensils before application. Wash teats of animals before milking.

- When used in indoor food handling/ processing facilities: Do not make space spray applications when facility is in operation. During space spray applications, cover or remove exposed food. During space spray applications, cover food processing surfaces or clean after treatment with a suitable detergent, rinse with potable water before use. Do not apply more than 1 time per day.
- For food crops growing outdoors or in greenhouses in agricultural setting: Do not apply more than 10 times per season. Do not re-apply within 3 days except under extreme pest pressure. In case of extreme pest pressure, do not re-apply within 24 hours. Do not harvest until spray has dried. Do not apply to cotton within 14 days of seed harvest.
- For post-harvest applications to vegetables, fruits, nuts and other commodities: Do not re-apply within 7 days. Do not apply more than 10 times to sweet potatoes.
- For post-harvest applications to stored grains and seed: Do not re-apply within 30 days.
- For greenhouse grown ornamental, flowering and foliage plants: Do not apply more than 1 time per day.
- Remove pets, birds and cover fish aquariums before application.

ENTRY RESTRICTIONS:

For Surface Spray Use: Except when applying to livestock and pets and when used as a wide area mosquito adulticide, do not enter or allow others to enter treated areas until sprays have dried.
For Space Spray Use: Do not enter or allow others to enter until vapors, mists, and aerosols have dispersed, and the treated area has been thoroughly ventilated.

Evergreen, McLaughlin Gormley King Company, MGK and Premium Pyrethrum are registered trademarks of McLaughlin Gormley King Company. All other marks are property of their respective holders. Manufactured by McLaughlin Gormley King Company®. ©2010 McLaughlin Gormley King Company. All rights reserved. 8810 Tenth Avenue North, Minneapolis, MN 55427

2.5M-0713

