

Supplemental Labeling

RESTRICTED USE PESTICIDE

DUE TO ACUTE INHALATION TOXICITY OF SULFURYL FLUORIDE

For sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification. An applicator certified by the state must be present on site at all times during introduction of fumigant, reentry prior to aeration, and initiation of the aeration procedure.

ProFume®

EPA Reg. No. 1015-79

**24(c) Special Local Need Registration SLN PA-070005
(For Distribution and Use Only Within the State of Pennsylvania)**

Fumigation of Cocoa Bean for Control of Postharvest Insect Pests

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the product container before applying. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label attached to the product container.

Directions for Use

This Section 24(c) Special Local Needs label permits the use of ProFume® gas fumigant on cocoa bean for control of postharvest insects. Follow all use directions, requirements and precautions on the label attached to the product container and included in the ProFume Applicator's Manual.

Specific Use Restrictions:

- Do not exceed a total cumulative dosage of 3750 g-h/m³ (oz-hr/1000 ft³).
- **Maximum single fumigation dosage:** 1500 g-h/m³ (oz-hr/1000 ft³).
- The aeration process must be completed prior to subsequent fumigation. Following each aeration period, the fumigator must test the breathing spaces in the space or structure to make certain that the concentration of ProFume is ≤ 1 ppm before allowing re-occupation of the structure.

®Trademark of Douglas Products and Packaging Company

Douglas Products and Packaging Company
1550 East Old 210 Hwy
Liberty, MO 64068-9459